



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

to the sum of two squares, I have found the following complete sets of "eggs" for each nest.

In nest 1, where  $n=1$  and  $N=1$ : 1, 2, 2. PROOF.— $1^2+2^2+2^2=(3^1)^2=9$ .

In nest 2, where  $n=2$  and  $N=2$ : 1, 4, 8; 7, 4, 4.

In nest 3, where  $n=3$  and  $N=5$ : 7, 2, 26; 7, 14, 22; 23, 2, 14; 23, 10, 10; 25, 2, 10.

In nest 4, where  $n=4$  and  $N=14$ :

1, 28, 76; 1, 44, 68; 17, 56, 56; 23, 16, 76; 23, 44, 64; 41, 16, 68; 41, 28, 64; 47, 16, 64; 49, 8, 64; 49, 32, 56; 55, 20, 56; 55, 40, 44; 65, 20, 44; 79, 8, 16.

In nest 5, where  $n=5$  and  $N=41$ :

1, 22, 242; 17, 14, 242; 17, 46, 238; 17, 106, 218; 17, 134, 202; 31, 38, 238; 31, 158, 182; 47, 14, 238; 47, 154, 182; 49, 2, 238; 49, 158, 178; 95, 118, 190; 95, 50, 218; 95, 130, 182; 97, 46, 218; 97, 62, 214; 97, 94, 202; 97, 134, 178; 113, 22, 214; 113, 62, 206; 113, 74, 202; 113, 146, 158; 127, 22, 206; 127, 46, 202; 127, 106, 178; 127, 134, 158; 143, 50, 190; 143, 74, 182; 143, 122, 154; 145, 70, 182; 161, 2, 182; 161, 38, 178; 193, 22, 146; 193, 62, 134; 193, 70, 130; 209, 22, 122; 209, 38, 118; 223, 22, 94; 223, 62, 74; 239, 22, 38; 241, 22, 22.

In nest 6, where  $n=6$  and  $N=122$ :

17, 136, 716; 17, 436, 584; 49, 236, 668; 49, 224, 692; 49, 128, 716; 49, 496, 532; 55, 196, 700; 73, 116, 716; 73, 44, 724; 79, 32, 724; 79, 172, 704; 79, 112, 716; 79, 308, 656; 79, 340, 640; 79, 460, 560; 89, 224, 688; 89, 416, 592; 103, 376, 616; 103, 424, 584; 119, 68, 716; 119, 196, 692; 119, 436, 572; 119, 484, 532; 137, 4, 716; 137, 236, 676; 143, 124, 704; 143, 284, 656; 175, 196, 680; 175, 104, 700; 185, 40, 704; 185, 296, 640; 199, 136, 688; 199, 248, 656; 199, 304, 632; 199, 376, 592; 239, 308, 616; 241, 4, 688; 241, 416, 548; 241, 128, 676; 241, 464, 508; 247, 116, 676; 247, 236, 644; 271, 32, 676; 271, 220, 640; 271, 208, 644; 271, 380, 560; 313, 56, 656; 313, 304, 584; 329, 92, 644; 329, 460, 460; 337, 56, 644; 337, 196, 616; 359, 56, 632; 359, 152, 616; 359, 248, 584; 359, 424, 472; 367, 236, 584; 367, 296, 556; 401, 172, 584; 401, 248, 556; 401, 296, 532; 401, 364, 488; 409, 376, 472; 409, 152, 584; 431, 68, 584; 431, 136, 572; 431, 296, 508; 431, 376, 452; 439, 172, 556; 439, 196, 548; 439, 236, 532; 439, 284, 508; 455, 104, 560; 455, 280, 496; 457, 116, 556; 457, 364, 436; 473, 304, 464; 497, 196, 496; 497, 224, 484; 521, 44, 508; 521, 100, 500; 521, 220, 460; 521, 236, 452; 521, 340, 380; 527, 196, 464; 527, 284, 516; 529, 40, 500; 529, 116, 488; 529, 200, 460; 529, 268, 424; 529, 288, 436; 529, 332, 376; 551, 112, 464; 551, 304, 368; 583, 136, 416; 583, 224, 376; 593, 4, 424; 593, 196, 376; 623, 44, 376; 623, 104, 364; 623, 236, 296; 625, 196, 320; 625, 220, 304; 631, 28, 364; 631, 196, 308; 649, 4, 332; 649, 124, 308; 649, 172, 284; 649, 196, 268; 655, 4, 320; 655, 100, 304; 655, 40, 296; 655, 104, 280; 689, 32, 236; 689, 116, 208; 695, 4, 220; 695, 100, 196; 719, 32, 116; 719, 44, 112; 721, 28, 104; 721, 40, 100; 721, 56, 92.

In order to obtain the multiple sets in the several nests, we multiply the prime sets of all the preceding nests by the necessary power of 3.

The multiple sets in nest 4 are  $3^3$  times the prime set of nest 1,  $3^2$  times the prime sets of nest 2, and 3 times those of nest 3, making 8 in all.

Also solved by CHARLES C. CROSS.

## AVERAGE AND PROBABILITY.

56. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics in Drury College, Springfield, Mo.


Find the chance that the center of gravity of a triangle lies inside the triangle formed by three points taken at random within the triangle. [From *Williamson's Integral Calculus*.]

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let  $ABC$  be the given triangle,  $D, E, F$  the mid-points of the sides, and  $O$  the center of gravity. Also let  $DG=x$ ,  $DS, DH, DJ, FK, FL, FN, ER$ , or  $EI=y$ .

$$\text{Now } FM = BM - \frac{1}{2}c = \frac{c(a-2x)}{2(a+6x)}.$$

When the first point is on  $OG$  and the second on  $OH, OS, OJ, OK, OL, ON, OR$ , or  $OI$ , respectively, then the third point must fall on



$$MON = \frac{\Delta}{3} \left( \frac{4x}{a+6x} - \frac{4y}{a+6y} \right), \quad MOL = \frac{\Delta}{3} \left( \frac{4y}{a+6y} - \frac{4x}{a+6x} \right),$$

$$MARO = \frac{\Delta}{3} \left( \frac{4x}{a+6x} + \frac{4y}{a+6y} \right), \quad MAIO = \frac{\Delta}{3} \left( \frac{4x}{a+6x} + 1 - \frac{4y}{c+6y} \right),$$

$$MACSO = \frac{\Delta}{3} \left( \frac{4x}{a+6x} + 1 + \frac{4y}{c+6y} \right), \quad MBHO = \frac{\Delta}{3} \left( 2 - \frac{4x}{a+6x} - \frac{4y}{c+6y} \right),$$

$$MOJB = \frac{\Delta}{3} \left( \frac{4y}{b+6y} + 1 - \frac{4x}{a+6x} \right), \quad MOK = \frac{\Delta}{3} \left( 1 - \frac{4y}{b+6y} - \frac{4x}{a+6x} \right),$$

respectively.

The chance that the first is on  $OG$  is  $dx/3a$ ; that the second is on  $OH, OS$ , or  $OJ$  is  $dy/3a$ ; that it is on  $OK, OL$ , or  $ON$  is  $dy/3c$ ; that it is on  $OR$ , or  $OI$  is  $dy/3b$ .

$$\begin{aligned} \therefore p &= \frac{1}{27a^2} \left[ \int_0^{\frac{1}{2}a} \left\{ \int_0^x \left( \frac{4x}{a+6x} - \frac{4y}{a+6y} \right) dy + \int_x^{\frac{1}{2}a} \left( \frac{4y}{a+6y} - \frac{4x}{a+6x} \right) dy \right\} dx \right. \\ &+ \int_0^{\frac{1}{2}a} \int_0^{\frac{1}{2}a} \left( \frac{4x}{a+6x} + \frac{4y}{a+6y} \right) dx dy \Big] + \frac{1}{27ac} \left[ \int_0^{\frac{1}{2}a} \int_0^{\frac{1}{2}c} \left( \frac{4x}{a+6x} + 1 - \frac{4y}{c+6y} \right) dx dy \right. \\ &+ \int_0^{\frac{1}{2}a} \left\{ \int_0^{FM} \left( \frac{4x}{a+6x} + 1 + \frac{4y}{c+6y} \right) dy + \int_{FM}^{\frac{1}{2}c} \left( 2 - \frac{4x}{a+6x} - \frac{4y}{c+6y} \right) dy \right\} dx \Big] \\ &+ \frac{1}{27ab} \left[ \int_0^{\frac{1}{2}a} \int_0^{\frac{1}{2}b} \left( \frac{4y}{b+6y} + 1 - \frac{4x}{a+6x} \right) dx dy + \int_0^{\frac{1}{2}a} \int_0^{\frac{1}{2}b} \left( 1 - \frac{4y}{b+6y} - \frac{4x}{a+6x} \right) dx dy \right] \\ \therefore p &= \frac{1}{27a} \int_0^{\frac{1}{2}a} \left[ \frac{1}{2} + \frac{a-34x}{6(a+6x)} + \frac{4x(a-2x)}{(a+6x)^2} + \frac{4}{9} \log \left( \frac{a+6x}{a} \right) - \frac{2}{9} \log 4 \right] dx \\ &= \frac{1}{27} \left( \frac{1}{3} + \frac{5}{9} \log 4 \right). \end{aligned}$$

Since the first point can fall in any of the six portions into which the medians divide the triangle, the required probability is  $P=6p$ .

$$\therefore P = \frac{6}{2^7} \left( \frac{1}{3} + \frac{5}{9} \log 4 \right) = \frac{1}{2^7} \left( 2 + \frac{10}{9} \log 4 \right) = \frac{2}{2^7} (1 + \frac{5}{9} \log 2).$$


91. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Six points  $A, B, C, D, E, F$ , are taken at random on the surface of a sphere. Find the chance that the plane through  $A, B, C$  intersects the plane through  $D, E, F$  within the sphere.

Solution by the PROPOSER.

Let  $AI, DJ$  be the diameters of the sections of the sphere made by the planes through  $A, B, C$  and  $D, E, F$ ;  $M$  and  $N$  their centers,  $O$  the center of the sphere,  $OP$  a line such that  $AB$  is parallel to the plane  $MOP$ ,  $OQ$  a line such that  $DE$  is parallel to the plane  $NOQ$ . Draw  $AG, DH$  perpendicular to  $AI, DJ$ , respectively.

Let  $AO=r$ ,  $\angle AOM=\theta$ ,  $\angle DON=\varphi$ ,  $\angle GAC=\psi$ ,  $\angle GAB=\delta$ ,  $\angle HDF=\mu$ ,  $\angle HDE=\nu$ ,  $\angle MOP=\lambda$ , the angle the plane  $MOP$  makes with some fixed plane through  $OP=\rho$ ,  $\angle NOQ=\gamma$ , and the diedral angle  $MOQN=\eta$ .



An element of surface at  $A$  is  $4\pi r^2 \sin \theta d\theta$ ; at  $D$ ,  $4\pi r^2 \sin \varphi d\varphi$ ; at  $C$ ,  $4r^2 \sin \theta \sin \psi d\psi d\lambda$ ; at  $B$ ,  $4r^2 \sin \theta \sin(\psi - \delta) \sin \lambda \sin \delta d\delta d\rho$ ; at  $F$ ,  $4r^2 \sin \varphi \sin \mu d\mu d\gamma$ ; at  $E$ ,  $4r^2 \sin \varphi \sin(\mu - \nu) \sin \gamma \sin \nu d\nu d\eta$ .

The limits of  $\theta$  are 0 and  $\frac{1}{2}\pi$ ; of  $\varphi$ , 0 and  $\frac{1}{2}\pi$ ; of  $\psi$ , 0 and  $\pi$ ; of  $\delta$ , 0 and  $\psi$ ; of  $\mu$ , 0 and  $\pi$ ; of  $\nu$ , 0 and  $\mu$ ; of  $\lambda$ , 0 and  $\pi$ ; of  $\gamma$ ,  $\pm(\theta - \varphi)$  and  $\theta + \varphi$  (the double sign being taken  $+$  when  $\theta > \varphi$ , and  $-$  when  $\theta < \varphi$ ); of  $\rho$ , 0 and  $2\pi$ ; of  $\eta$ , 0 and  $2\pi$ .

Since the whole number of ways six points can be taken in the surface of the sphere is  $(4\pi r^2)^6$  the required chance is

$$\begin{aligned} p &= \frac{1}{(4\pi r^2)^6} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^\pi \int_0^\pi \int_0^\psi \int_0^\mu \int_0^\pi \int_{\pm(\theta-\varphi)}^{\theta+\varphi} \int_0^{2\pi} \int_0^{2\pi} 4\pi r^2 \sin \theta. d\theta 4\pi r^2 \\ &\quad \sin \varphi d\varphi. 4r^2 \sin \theta \sin \psi d\psi d\lambda. 4r^2 \sin \varphi \sin \mu d\mu d\gamma. 4r^2 \sin \theta \sin(\psi - \delta) \sin \lambda \sin \delta d\delta d\rho \\ &\quad \times 4r^2 \sin \varphi \sin(\mu - \nu) \sin \gamma \sin \nu d\nu d\eta \\ &= \frac{2}{\pi^3} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^\pi \int_0^\pi \int_0^\psi \int_0^\mu \int_0^\pi \int_{\pm(\theta-\varphi)}^{\theta+\varphi} \int_0^{2\pi} \sin^3 \theta \sin^3 \varphi \sin \psi \sin \mu \sin(\psi - \delta) \\ &\quad \sin(\mu - \nu) \sin \delta \sin \nu \sin \lambda \sin \gamma d\theta d\varphi d\psi d\mu d\delta d\nu d\gamma d\rho \\ &= \frac{4}{\pi^2} \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} \int_0^\pi \int_0^\pi \int_0^\psi \int_0^\mu \int_0^\pi \int_{\pm(\theta-\varphi)}^{\theta+\varphi} \sin^3 \theta \sin^3 \varphi \sin \psi \sin \mu \sin(\psi - \delta) \end{aligned}$$